

## Cholesterol (CHOL)

Source:

Cerba Lancet Africa

View our website:

[cerbalancetafrica.com](http://cerbalancetafrica.com)

Date & Time of Export:

09 January 2026 00:50



### TEST OVERVIEW

<b>Test Name</b>	Cholesterol
<b>Test Code</b>	CHOL
<b>Short Description</b>	CHOL

### OVERVIEW

<b>Test Name</b>	Cholesterol
<b>Test Code</b>	CHOL
<b>Category</b>	Biochemistry
<b>TAT</b>	Main Lab: 6 Hour(s) Family Site: <8hrs
<b>Specimen(s)</b>	1 x Venous blood - 5 mL Tube - Gold - SST-Serum Separator Tube

### SPECIMEN(S)

#### SST-Serum Separator Tube

<b>Specimen Type</b>	SST-Serum Separator Tube
<b>Specimen Format</b>	Tube
<b>Specimen Colour</b>	Gold
<b>Specimen Volume</b>	5 mL
<b>Sampling Order</b>	2
<b>Origin</b>	Venous blood
<b>Collection time after baseline</b>	-
<b>Transport Temperature</b>	15-25°C
<b>Accepted Other Specimens</b>	Lithium Heparin Plasma Sodium Heparin Plasma Serum

**TAT**

EDTA Plasma

Main Lab: 6 Hour(s)  
Family Site: <8hrs**Test Stability**Room Temp: 7 Day(s)  
2–8°C: 7 Day(s)**PATIENT INFORMATION****Clinical Information Required**

Fasting patient

**Patient Collection Notes**

In the absence of known hypertriglyceridaemia, a lipid profile may be taken in non-fasting patients. If hypertriglyceridaemia is discovered, a control sample should be prescribed and taken after a 12-hour fast. In all cases, the patient may drink water.

**COMMENTS & NOTES****LOINC Code**

647-2, 14647-2,

**Outwork**

No