Rickettsial screen (IgG+IgM) (RICKET_SCR)

Source: Cerba Lancet Africa

View our website: cerbalancetafrica.com

Date & Time of Export: 23 November 2026 04:16

Click here to view the results online

TEST OVERVIEW

Test Name	Rickettsial screen (IgG+IgM)
Test Code	RICKET_SCR
Short Description	Rickettsial screen (IgG+IgM)

OVERVIEW

Test Name	Rickettsial screen (IgG+IgM)
Test Code	RICKET_SCR
Category	Serology
TAT	Main Lab: 2 Day(s) Family Site: 2 Days
Specimen(s)	1 x Venous blood - 5 mL Tube - Gold - SST

SPECIMEN(S)

SST

Specimen Type	SST
Specimen Format	Tube
Specimen Colour	Gold
Specimen Volume	5 mL
Sampling Order	2
Origin	Venous blood
Collection time after baseline	-
Transport Temperature	15-25°C
Accepted Other Specimens	Serum
TAT	Main Lab: 2 Day(s)

	Family Site: 2 Days
Test Stability	Room Temp: 8 Hour(s) 2–8°C: 7 Day(s)

CLINICAL INFORMATION

Rickettsial screen (IgG+IgM)

Methodology	-
Specimen Type	SST
Delay before pre-treatment	4
Transport Temperature	15-25°C
Transport Stability at room temp	8 Hours
Transport Stability at 2–8°C	7 Day
Haemolysis interference	No

PATIENT INFORMATION

Clinical Information Required	-
Patient Collection Notes	-

COMMENTS & NOTES

LOINC Code	94-7, 7994-7
Outwork	Yes